

Table 1.13A Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Numbers in Thousands, 2011 and 2012

Drug	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
ILLCIT DRUGS¹	33,552 ^b	37,011	9,844 ^b	11,655	2,888	3,068	2,490 ^b	3,291	4,466 ^b	5,297	23,707 ^a	25,356
Marijuana and Hashish	26,186 ^b	28,150	7,127 ^b	8,528	2,052	2,298	1,821 ^b	2,335	3,253 ^b	3,895	19,059	19,622
Cocaine	3,628 ^b	4,491	1,210 ^b	1,740	434	504	261 ^b	542	516	694	2,418	2,751
Crack	597 ^a	902	291	460	108	174	71	129	112	157	306	442
Heroin	566	638	366	403	144	181	84	85	137	137	201	236
Hallucinogens	3,421	3,753	1,084	1,216	378	369	264	321	443	526	2,336	2,537
LSD	721	905	235	298	86	112	49	70	100	116	486	607
PCP	80	124	37	34	10	24	11	6	16	4	43	90
Ecstasy	2,003	2,322	675	783	233	225	156	213	285	345	1,328	1,539
Inhalants	1,041	1,046	390	458	114	163	119	155	157	140	651	587
Nonmedical Use of Psychotherapeutics ^{2,3}	12,908 ^b	15,025	4,764 ^b	5,811	1,517	1,758	1,230 ^a	1,605	2,016 ^a	2,448	8,144 ^b	9,213
Pain Relievers	9,659 ^b	11,157	3,586 ^a	4,181	1,162	1,286	1,024	1,136	1,400 ^a	1,759	6,073 ^a	6,976
OxyContin [®]	1,404	1,290	667	743	198	282	204	203	265	258	738 ^a	547
Tranquilizers	4,635 ^b	5,621	1,948 ^b	2,747	676	873	491 ^a	747	781 ^a	1,127	2,686	2,874
Stimulants ³	2,401 ^b	2,992	1,043 ^a	1,351	393	480	231 ^a	388	419	484	1,358	1,641
Methamphetamine ³	928	1,073	507	577	197	233	127	138	183	207	420	496
Sedatives	442	509	252	277	123	108	73	92	57	78	190	231
ILLCIT DRUGS OTHER THAN MARIJUANA¹	16,395 ^b	18,970	5,724 ^b	6,937	1,811	1,954	1,459 ^b	1,967	2,454 ^a	3,016	10,671 ^b	12,033

*Low precision; no estimate reported.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.13B Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Percentages, 2011 and 2012

Drug	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
ILLCIT DRUGS¹	14.4 ^b	15.7	23.8 ^b	26.7	32.1	32.1	24.1	27.6	20.3 ^b	23.8	12.4 ^a	13.2
Marijuana and Hashish	11.3 ^a	12.0	17.2 ^b	19.5	22.8	24.0	17.6	19.6	14.8 ^b	17.5	10.0	10.3
Cocaine	1.6 ^b	1.9	2.9 ^b	4.0	4.8	5.3	2.5 ^b	4.5	2.3	3.1	1.3	1.4
Crack	0.3 ^a	0.4	0.7	1.1	1.2	1.8	0.7	1.1	0.5	0.7	0.2	0.2
Heroin	0.2	0.3	0.9	0.9	1.6	1.9	0.8	0.7	0.6	0.6	0.1	0.1
Hallucinogens	1.5	1.6	2.6	2.8	4.2	3.9	2.6	2.7	2.0	2.4	1.2	1.3
LSD	0.3	0.4	0.6	0.7	1.0	1.2	0.5	0.6	0.5	0.5	0.3	0.3
PCP	0.0	0.1	0.1	0.1	0.1	0.3	0.1	0.1	0.1	0.0	0.0	0.0
Ecstasy	0.9	1.0	1.6	1.8	2.6	2.4	1.5	1.8	1.3	1.6	0.7	0.8
Inhalants	0.4	0.4	0.9	1.0	1.3	1.7	1.1	1.3	0.7	0.6	0.3	0.3
Nonmedical Use of Psychotherapeutics ^{2,3}	5.5 ^b	6.4	11.5 ^a	13.3	16.8	18.4	11.9	13.5	9.1 ^a	11.0	4.3 ^b	4.8
Pain Relievers	4.2 ^b	4.7	8.7	9.6	12.9	13.4	9.9	9.5	6.4 ^a	7.9	3.2 ^a	3.6
OxyContin [®]	0.6	0.5	1.6	1.7	2.2	2.9	2.0	1.7	1.2	1.2	0.4 ^a	0.3
Tranquilizers	2.0 ^b	2.4	4.7 ^b	6.3	7.5	9.1	4.8	6.3	3.5 ^a	5.1	1.4	1.5
Stimulants ³	1.0 ^a	1.3	2.5	3.1	4.4	5.0	2.2	3.2	1.9	2.2	0.7	0.9
Methamphetamine ³	0.4	0.5	1.2	1.3	2.2	2.4	1.2	1.2	0.8	0.9	0.2	0.3
Sedatives	0.2	0.2	0.6	0.6	1.4	1.1	0.7	0.8	0.3	0.3	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	7.0 ^b	8.1	13.8 ^b	15.9	20.1	20.4	14.1	16.5	11.1 ^a	13.6	5.6 ^b	6.3

*Low precision; no estimate reported.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.14A Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Numbers in Thousands, 2011 and 2012

Substance	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
PAST YEAR												
Tobacco Products ¹	77,656	79,029	17,909 ^a	19,284	4,744	4,804	4,667 ^a	5,423	8,498	9,057	59,747	59,745
Cigarettes	63,839	64,854	15,851 ^a	17,079	4,369	4,349	4,131 ^b	4,981	7,351	7,749	47,988	47,775
Smokeless Tobacco	10,621	11,069	1,715	2,074	398	547	312 ^a	501	1,005	1,026	8,905	8,995
Cigars	23,040	23,964	4,648 ^a	5,341	1,263	1,385	1,152 ^a	1,488	2,233	2,468	18,392	18,622
Alcohol	163,482 ^b	167,018	28,927 ^b	31,270	6,640	6,981	7,335 ^b	8,686	14,952	15,603	134,555	135,748
PAST MONTH												
Tobacco Products ¹	65,735	67,345	15,571 ^a	16,815	4,259	4,227	3,995 ^a	4,728	7,317	7,860	50,164	50,530
Cigarettes	54,880	55,890	13,880 ^a	15,019	3,950	3,821	3,579 ^a	4,319	6,351	6,878	41,000	40,872
Smokeless Tobacco	7,726 ^a	8,521	1,180	1,434	242	317	187 ^a	334	751	784	6,546	7,087
Cigars	12,007	12,772	2,516	2,879	643	769	579 ^a	933	1,294	1,177	9,491	9,893
Pipe Tobacco	1,955	2,365	534	720	140	174	197	214	197	332	1,421	1,645
Alcohol	130,059	132,285	22,029 ^b	24,045	4,840	5,218	5,392 ^b	6,579	11,797	12,247	108,030	108,240
Binge Alcohol Use ²	56,491	57,886	10,569 ^b	12,053	2,552	2,822	2,652 ^a	3,252	5,365	5,979	45,922	45,833
Heavy Alcohol Use ²	15,486 ^a	16,695	3,314 ^a	4,036	867	1,022	822	1,036	1,625	1,978	12,171	12,659

*Low precision; no estimate reported.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.14B Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Percentages, 2011 and 2012

Substance	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
PAST YEAR												
Tobacco Products ¹	33.4	33.6	43.3	44.1	52.6	50.2	45.2	45.5	38.6	40.8	31.2	31.2
Cigarettes	27.4	27.6	38.3	39.1	48.5	45.4	40.0	41.8	33.4	34.9	25.1	25.0
Smokeless Tobacco	4.6	4.7	4.1	4.7	4.4	5.7	3.0	4.2	4.6	4.6	4.7	4.7
Cigars	9.9	10.2	11.2	12.2	14.0	14.5	11.2	12.5	10.1	11.1	9.6	9.7
Alcohol	70.3	71.0	69.9	71.5	73.7	73.0	71.0	72.8	67.8	70.2	70.4	70.9
PAST MONTH												
Tobacco Products ¹	28.3	28.6	37.6	38.5	47.3	44.2	38.7	39.6	33.2	35.4	26.2	26.4
Cigarettes	23.6	23.8	33.5	34.4	43.8	39.9	34.6	36.2	28.8	31.0	21.4	21.4
Smokeless Tobacco	3.3	3.6	2.9	3.3	2.7	3.3	1.8	2.8	3.4	3.5	3.4	3.7
Cigars	5.2	5.4	6.1	6.6	7.1	8.0	5.6	7.8	5.9	5.3	5.0	5.2
Pipe Tobacco	0.8	1.0	1.3	1.6	1.6	1.8	1.9	1.8	0.9	1.5	0.7	0.9
Alcohol	55.9	56.3	53.2	55.0	53.7	54.5	52.2	55.2	53.5	55.1	56.5	56.5
Binge Alcohol Use ²	24.3	24.6	25.5 ^a	27.6	28.3	29.5	25.7	27.3	24.3	26.9	24.0	23.9
Heavy Alcohol Use ²	6.7	7.1	8.0	9.2	9.6	10.7	8.0	8.7	7.4	8.9	6.4	6.6

*Low precision; no estimate reported.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.